



Review

Combined extracorporeal shockwave therapy and exercise for the treatment of tendinopathy: A narrative review

Ian Burton

NHS Grampian, 172 Market Street, Aberdeen, AB11 5PP, United Kingdom

ARTICLE INFO

Keywords:

Exercise
Muscle stretching exercises
High-energy shock waves
Tendinopathy
Resistance training
Extracorporeal shockwave therapy

ABSTRACT

Tendinopathy is a chronic degenerative musculoskeletal disorder that is common in both athletes and the general population. Exercise and extracorporeal shockwave therapy (ESWT) is among the most common treatments used to mediate tendon healing and regeneration. The review presents the current understanding of mechanisms of action of ESWT and exercise in isolation and briefly synthesises evidence of their effectiveness for various tendinopathies. The central purpose of the review is to synthesise research findings investigating the combination of ESWT and exercise for five common tendinopathies (plantar heel pain, rotator cuff, lateral elbow, Achilles, and patellar tendinopathy) and provide recommendations on clinical applicability. Collectively, the available evidence indicates that ESWT combined with exercise in the form of eccentric training, tissue-specific stretching, or heavy slow resistance training are effective for specific tendinopathies and can therefore be recommended in treatment. Whilst there are at present a limited number of studies investigating combined ESWT and exercise approaches, there is evidence to suggest that the combination improves outcomes in the treatment of plantar heel pain, Achilles, lateral elbow, and rotator cuff tendinopathy. However, despite overall positive outcomes in patellar tendinopathy, the combined treatment has not been shown to offer additional benefit over eccentric exercise alone.

Introduction

Tendinopathy is a degenerative tendon disorder associated with chronic pain, swelling and impaired physical function.^{1,2} Tendons are susceptible to injury as they undergo high forces during loading activities, receive little vascular blood supply, have low elasticity, and decreased metabolism.^{3,4} Tendinopathy is thought to be caused by repetitive tendon microtrauma, and subsequently failed healing responses, characterised by neovascularization, incidence of calcium deposits or calcification, and increased tendon thickness or swelling.⁵ Although inflammatory cells are typically present in tendinopathy, tendon cell degeneration rather than inflammation is considered the pathological hallmark of the disorder, leading to the adoption of the term tendinopathy rather than the previously used diagnosis of ‘tendinitis’.⁶ Recent data from Denmark and the Netherlands on lower limb tendinopathies estimate incidence and prevalence ranging from 7.0 to 11.8 and 10.5 to 16.6 per 1000 people, respectively.^{7,8} Athletes exposed to high musculoskeletal loads are particularly susceptible to developing tendinopathy, particularly if these loads are experienced frequently, which can lead to repeated tendon stress and overloading.⁹ Prevalence of rotator cuff tendinopathy has been reported to be 30% in volleyball players,¹⁰ and 13%

of climbers were reported to have lateral elbow tendinopathy in one study.¹¹ Studies on patellar tendinopathy have reported a prevalence of 17% in elite basketball players¹² and 6% in adolescent athletes.¹³ Achilles tendinopathy prevalence has been reported to be 135% in adult athletes¹⁴ and 8% in adolescent runners.¹⁵

Tendinopathies are considered to have a multifactorial pathogenesis resulting from a range of extrinsic and intrinsic factors.¹⁶ Pain experienced from tendinopathy is not always associated with tissue damage, suggesting other neurophysiological or psychological mechanisms are involved.^{17,18} Central nervous system hypersensitivity and the role of pain central sensitisation has recently emerged as a potential contributing factor.¹⁹ Identified risk factors for tendinopathy include altered biomechanics,²⁰ rheumatological and metabolic disorders such as diabetes,²¹ excess adiposity,²² and medications such as fluoroquinolones.²³ A range of approaches are used in tendinopathy rehabilitation including anti-inflammatory medications, corticosteroid injections, low-level-laser therapy, ultrasound, platelet-rich-plasma injections, prolotherapy, glycerol trinitrate patches, manual therapy, and exercise.²⁴ Exercise is the most commonly recommended treatment option, with specific exercise targeting the injured tendons the most common exercise approach for tendinopathy.²⁵ In the last decade, there has been a growing body of research investigating the effectiveness of Extracorporeal shockwave

E-mail address: ianburton_10@hotmail.co.uk.

<https://doi.org/10.1016/j.smhs.2021.11.002>

Received 11 September 2021; Received in revised form 5 November 2021; Accepted 5 November 2021

Available online 11 November 2021

2666-3376/© 2021 Chengdu Sport University. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd.

Abbreviations

ESWT	Extracorporeal shockwave therapy
F-ESWT	Focused extracorporeal shockwave therapy
LLLT	Low level laser therapy
PHP	Plantar heel pain
PFSS	Plantar fascia specific stretching
R-ESWT	Radial extracorporeal shockwave therapy
RCTs	Randomised controlled trials

therapy (ESWT) as a tendinopathy treatment.^{26,27} Musculoskeletal conditions including those involving tendons, ligaments, muscles, joints, and bones can be treated effectively with ESWT.²⁸ Due to its non-invasive nature, dearth of side effects, and patient acceptability, ESWT offers a therapeutic rehabilitation method when other conservative treatments are ineffective.²⁹ Although there have been many reviews presenting the evidence on ESWT and exercise in isolation for tendinopathies, there are currently none which have examined a combined approach. The purpose of this review is to synthesize the currently available research on exercise and ESWT for five tendinopathies (plantar heel pain, rotator cuff, lateral elbow, patellar, and Achilles), with a focus on the evidence for combined ESWT and exercise interventions for these tendinopathies. The review will begin with an overview of mechanisms of action for ESWT and exercise in tendinopathy and include a summary of the effectiveness of each in isolation, and finally an overview of studies that have combined ESWT and exercise, concluding with clinical recommendations.

Methodology

To conduct this narrative review, a systematic literature search was conducted using Medline/PubMed database to identify studies up to March 2020 using a combination of the following search terms: extracorporeal shockwave therapy, shock wave, shockwave therapy, shockwave, ESWT, tendon, tendinopathy, musculoskeletal, plantar, Achilles, patellar, rotator cuff, elbow, epicondylitis, exercise, strengthening, eccentric, resistance training, physical therapy, and physiotherapy. Relevant articles were retrieved and used to identify additional sources by cross-referencing and manually checking reference lists. The findings of relevant research investigating combined ESWT and exercise interventions are presented in [Table 1](#).

Exercise mechanisms in tendinopathy

Heavy slow resistance training and eccentric strengthening interventions have been shown to be the most evidence-based exercise approach for tendinopathy, having good outcomes for various tendinopathies such as plantar heel pain³⁰ rotator cuff,³¹ lateral elbow,³² Achilles,³³ and patellar tendinopathy.³⁴ The emerging evidence for exercise has resulted in investigations using eccentric training and heavy slow resistance training to treat tendinopathy through utilising the process of mechano-transduction.³⁵ The high forces and loads produced during slowly performed concentric and eccentric contractions may stimulate tendon remodelling and compliance by stimulating collagen synthesis and reducing neovascularization.³⁶ Through the process of mechano-transduction, exercise can influence cell homeostasis and therefore stimulate tendon regeneration.³⁷ Mechanical stimulus from exercise can upregulate cellular responses which alter structural adaptations in tendinopathy such as disorganized collagen architecture and increased water content in the extracellular matrix.³⁸ Structural changes in tendinopathy can alter the capacity of tendons to store and produce kinetic energy, affecting strength and functional performance.³⁹ Although loading forces applied to tendons during exercise may help stimulate remodelling of abnormal tendon structure, this may not be

Table 1

Studies investigating combined ESWT and exercise interventions for tendinopathies.

Study and design	Sample	Interventions	Outcome Measures	Results
Achilles tendinopathy				
Rompe 2009 RCT	n = 68	1: 12-week eccentric exercise protocol. 2: eccentric exercise plus R-ESWT for 3 sessions.	VISA-A, VAS.	Combination more effective for pain and function at 4 months, no difference between groups at 1 year.
Mansur 2017 Prospective cohort	n = 19	2 sessions of R-ESWT, and 12-week eccentric strengthening protocol.	VAS, AOFAS, VISA-A	Intervention considered effective for pain and function at 24 weeks.
Pavone 2016 Case series	n = 40	4 sessions of F-ESWT combined with eccentric exercises.	VAS, AOFAS	Combined intervention effective at 12 months and recommended as superior to eccentric exercise only.
Wheeler 2019 Case series	n = 39	3 sessions of R-ESWT and Home exercise: stretching, isometric and eccentric strengthening, core stability and proprioception exercises.	VISA-A	Significant improvements in insertional tendinopathy pain and function at 6 months but not for non-insertional tendinopathy.
Wheeler 2020 Prospective Cohort	n = 63	1: 3 sessions of ESWT. 2: High volume image guided injection. Both groups received home eccentric strengthening.	VAS, VISA-A, MOXFQ.	Significant improvements for pain and function in both groups, no significant differences between groups at 3 months.
Patellar tendinopathy				
Thijs et al. 2017 RCT	n = 52	1: 3 sessions of F-ESWT. 2: 3 sessions of sham ESWT. Both groups received eccentric squat exercises for 3 months	VISA-P	Both groups effective at 24 weeks, no additional effect of ESWT to eccentric exercises.
van der worp 2014 RCT	n = 43 (57 tendons)	1. 3 sessions of F-ESWT 2. 3 sessions of R-ESWT Both groups received eccentric training based on decline squats.	VISA-P	Both treatments effective at 14 weeks. No statistically significant differences in effectiveness between F-ESWT and R-ESWT.
van Rijn 2019 Secondary analysis from 3 RCTs	n = 138	1. ESWT 2. ESWT plus eccentric training 3. eccentric training 4. topical glyceryl trinitrate patch plus eccentric training 5. Placebo	VISA-P	Clinical improvements were significantly higher in the eccentric training and ESWT plus eccentric training groups compared to other groups.
Vetrano 2013 RCT	n = 46: athletes only.	1. 2 PRP injections. 2. 3 sessions of F-	VAS, VISA-P, Modified	PRP plus exercise had significantly better

(continued on next page)

Table 1 (continued)

Study and design	Sample	Interventions	Outcome Measures	Results
Rotator cuff tendinopathy	n = 30	ESWT	Blazina score.	improvement at 6 and 12 months.
		Both groups received stretching and strength exercise for 2 weeks		
		1: 3 sessions of F-ESWT. 2: F-ESWT plus 10 supervised sessions of isokinetic exercise.	VAS, CMS	Combined group had greater reduction of pain, and superior functional recovery and muscle endurance at 2 months compared with F-ESWT
Santamato 2016 RCT				
Kvalvaag 2018 RCT	n = 142	1. 4 sessions of Sham R-ESWT 2. 4 sessions of R-ESWT. Both groups received supervised exercise for 12-weeks.	SPADI	Both groups improved at 24 weeks and 1 year but there were no differences between the groups. Patients with calcification had a greater improvement at 24 weeks but not 1 year with R-ESWT plus exercise.
Carlisi 2018 Pilot RCT	n = 22	1. F-ESWT plus supervised eccentric training 2. F-ESWT only	p-NRS, DASH	Both groups improved pain and function at 9 weeks with no difference between groups. F-ESWT no benefit to eccentric exercise.
Lateral elbow tendinopathy				
Celik 2019 RCT	n = 43	1. 12 sessions of PBMT 2. 4 sessions of F-ESWT.	VAS, Strength, GRC	Improvements for VAS, elbow extension and shoulder flexion strength were superior in the PBMT group. Handgrip strength improved in both groups. More patients in PBMT group reported improvement with GRC. Both treatments effective with PBMT superior to ESWT.
		Both groups received home stretching and eccentric strengthening exercises.		
Eraslan 2018 RCT	n = 45	1: 15 sessions of physiotherapy with home exercise including stretching and eccentric strengthening 2: physiotherapy plus kinesio taping for 3 weeks. 3. physiotherapy	VAS, Grip strength, PRTEE	Pain and function improved in all groups. The kinesio taping and ESWT groups improved function better than the physiotherapy group.

Table 1 (continued)

Study and design	Sample	Interventions	Outcome Measures	Results
Testa 2020 Case series	n = 60	plus 3 sessions of ESWT 4 sessions of F-ESWT combined with eccentric exercises 4 times per week for one month.	VAS, PRTEE-1	Combined treatment effective for pain and function at 1, 6 and 12 months.
Plantar heel pain				
Akinoglu 2017 RCT	n = 54 females only.	1. Home exercise including PFSS for 4 weeks 2. 3 sessions of R-ESWT plus exercise 3. 7 sessions of US plus exercise	FFI, AOFAS.	Improved pain and function in all groups with US more effective than other groups. R-ESWT plus exercise superior to exercise alone.
Chew 2013 RCT	n = 53	1. Home gastrocnemius stretching and PFSS 2. ACP injection plus stretching 3. 2 sessions of ESWT plus stretching	VAS, AOFAS, VAS, plantar fascia thickness	ACP and ESWT plus stretching more effective for pain and function than stretching at 6 months. No significant difference between ACP and ESWT.
Cinar 2018 RCT	n = 66	1. Insoles for 3 months and stretching. 2. 3 sessions of R-ESWT, insoles and stretching. 3. 10 sessions LLLT, insoles and stretching	FFI-p, NRS-p.	All groups effective, LLLT and ESWT more effective than usual care, LLLT more effective than ESWT in reducing pain at 3 months.
Vahdatpour 2018 RCT	n = 80	1. 4 sessions of ESWT plus topical corticosteroid and home stretching 2. 4 sessions of ESWT plus Vaseline and stretching	VAS, RMS.	Both groups effective, combined group more effective than control at 1 month for pain and function, no difference at 3 months.
Takla 2019 RCT	n = 120	1. sham-PBMT plus stretching for 3 weeks 2. 9 sessions of PBMT plus stretching 3. 3 sessions of ESWT plus stretching 4. 3 sessions ESWT, 9 sessions of PBMT and stretching.	VAS, FFI-d, PPT.	Both ESWT and PBMT were effective for pain and function at 12 weeks. PBMT with ESWT was superior to ESWT and PBMT alone, and ESWT was superior to PBMT.
Grecco 2013 RCT	n = 40	1. 10 sessions of US plus home stretching 2. 3 sessions of R-ESWT plus home stretching.	VAS, PPT	Both treatments effective for pain and function. R-ESWT superior at 3 months, no difference between groups at 12 months.
Eslamian 2016 RCT	n = 40	1. 5 sessions of R-ESWT and home stretching 2. Single CSI and stretching	VAS, FFI	Both groups effective for pain and function at 8 weeks, ESWT superior to CSI.
Rompe 2015 RCT	n = 152, PHP 12 months	1. 3 sessions of ESWT plus PFSS 2. 3 sessions of ESWT	FFI, Morning pain	Combined group more effective for pain and function at 4

(continued on next page)

Table 1 (continued)

Study and design	Sample	Interventions	Outcome Measures	Results
Ulusoy 2017 RCT	n = 60, PHP 6 months or longer	1. 15 sessions of LLLT and home stretching 2. 15 sessions of US and home stretching 3. 3 sessions of ESWT and home stretching	VAS, AOFAS.	months, no difference between groups at 24 months. All groups effective for pain and function. LLLT and ESWT resulted in similar outcomes and were more successful than US.
Okur 2019 RCT	n = 83	1. 3 sessions of R-ESWT plus home stretching for 4 weeks 2. Custom orthotics plus home stretching for 4 weeks	VAS	Both groups effective in reducing pain with neither group superior at 24 nor 48 weeks.
Wheeler 2018 Case series	n = 35, PHP duration 24 months	3 sessions of R-ESWT and home exercise: stretching, foot and calf strengthening, balance exercises for 3 weeks.	VAS, FFI, MOXFQ	Intervention effective for pain and function at 3 months, but not in overall markers of health, anxiety/depression scores, or activity levels.

Abbreviations: AOFAS = American Orthopaedic Foot and Ankle Society; CMS = Constant-Murley score; CSI = Corticosteroid injection; DASH = disabilities of the arm, shoulder, and hand; ESWT = extracorporeal shockwave therapy; F-ESWT = focused extracorporeal shockwave therapy; FFI = Foot function index; GRC = Global Rating of Change; LLLT = low level laser therapy; MOXFQ = The Manchester-Oxford Foot Questionnaire; n = number; PBMT = photobiomodulation therapy; PRP = Platelet-rich plasma; PRTEE-1 = Patient Rated Tennis Elbow Evaluation Test; PHP = Plantar heel pain; p-NRS = pain numeric rating scale; PPT = Pain pressure threshold; RCT = randomised controlled trial; R-ESWT = radial extracorporeal shockwave therapy; RMS = Roles and Maudsley score; SPADI = The Shoulder Pain and Disability Index; US = ultrasound; VAS = Visual Analogue Scale; VISA-A = Victorian Institute of Sport Assessment-Achilles score; VISA-P = Victorian Institute of Sport Assessment-Patella score.

associated with increased collagen turnover, leading to uncertainties regarding healing mechanisms.⁴⁰ Although tendon structure can be altered in those under age 25 with loading exercise, adult tendons have reduced capacity for collagen turnover, therefore tendon changes do not adequately explain the clinical response to loading.^{41–43} Clinical improvement from exercise loading interventions may therefore be due to improved tendon mechanical properties, rather than altered tendon structure.⁴⁴

ESWT mechanisms in tendinopathy

In recent years, ESWT has been shown to have beneficial effects on musculoskeletal disorders and has emerged as an effective tendinopathy treatment.⁴⁵ There is now a growing body of evidence indicating the effectiveness of ESWT for both lower limb⁴⁶ and upper limb tendinopathies.⁴⁷ After first being introduced in urology to treat kidney stones, ESWT is now therapeutically applied to a broad range of medical and musculoskeletal disorders and is considered a significant innovation in regenerative medicine.⁴⁸ Two types of ESWT are used to treat musculoskeletal disorders, with focused-ESWT (F-ESWT) targeting deep smaller tissues, and radial-ESWT (R-ESWT) targeting larger superficial tissues.⁴⁹ Both types of ESWT generate high pressures which can induce cavitation in human tissues. The cavitation process involves rapid formation, expansion, and forceful collapse of vapor bubbles in liquids due to rapid

pressure changes.⁵⁰ Cavitation stimulates a plethora of regenerative biological responses in musculoskeletal tissues which activates proteins involved in chondroprotection,⁵¹ neovascularization,⁵² angiogenesis,⁵³ anti-inflammation,⁵⁴ anti-apoptosis,⁵⁵ immunomodulation,⁵⁶ neurophysiological and analgesic processes.⁵⁷ Healing processes are also activated, which induce proliferation, differentiation, and migration of various cells, including mesenchymal stem cells, stromal cells, endothelial cells, osteoblasts, fibroblasts, tenocytes, and enhancing collagen synthesis.⁴⁸ Several other tendon-specific responses which stimulate healing have been observed, including a reduction in pathological matrix constituents and inflammatory cytokines, upregulation of tendon cells, lubricin expression, anti-inflammatory cytokines, and collagen synthesis.⁵⁸

Combined treatment

In clinical practice, ESWT is rarely used in isolation and is often combined with specific exercises, aimed at loading damaged tendons to stimulate healing.⁵⁹ However, studies have generally not adequately addressed this, often comparing ESWT alone with placebo or usual care.⁶⁰ Recently a tendon pathology continuum model has been proposed, with tendinopathy presenting in three distinct stages, which would likely respond differently to ESWT and exercise.⁶¹ There is a need to consider comprehensive interventions depending on the type, stage, and characteristics of the tendinopathy.⁶² As monotherapy is rarely implemented clinically, there is a need for a comparison of more realistic treatment interventions.⁶³ Questions regarding combined ESWT and exercise effectiveness applied realistically as polytherapy in tendinopathy are unanswered.⁶⁰ There is increasing high-quality evidence for the clinical effectiveness of ESWT in tendinopathy, but systematic reviews still suggest a need for higher quality studies.⁶⁴ However, the remaining questions are not if ESWT works in isolation, but rather what treatment combinations are most effective for specific tendinopathies. Research needs to follow the clinical practice, where tendinopathy treatment is delivered as combined interventions.²⁶ A recent qualitative study in tendinopathy highlighted that patients felt ESWT alone was inadequate but acceptable as a non-invasive treatment combined with self-management exercise, which they regarded as more important.⁶⁵ Current evidence suggests that combined rather than single interventions should be recommended to achieve decreased pain, improved function, and patient satisfaction in tendinopathy treatment.^{66,67} Although combined treatments are recommended there is a dearth of randomised controlled trials (RCTs) assessing their effectiveness with or compared to other efficacious interventions for tendinopathies.⁶⁸ Recently studies have begun investigating combined ESWT with exercise for tendinopathies, with an overview of these studies and outcomes for each tendinopathy provided in the following sections.

Rotator cuff tendinopathy

Despite the aetiology of rotator cuff tendinopathy not being fully understood, a combination of extrinsic and intrinsic mechanisms is thought to be involved.¹⁶ Shoulder impingement is commonly involved, leading to tendon inflammation and eventual degeneration.⁶⁹ Symptoms include pain, weakness, and tenderness in the shoulder, especially when lifting overhead or abducting the arm beyond 90°.⁷⁰ Some patients can have long-term shoulder impairment, with 42% reporting symptoms after ten years.⁷¹ Calcific rotator cuff tendinopathy is a common cause of shoulder pain in those aged 30–60, with mechanical overloading a key predisposing factor.⁷² Calcium hydroxyapatite deposition characterizes calcification, with the supraspinatus tendon being implicated in 80% of cases.⁷³ Specific exercise strategies focussing on correcting kinematic deficits or eccentric only strengthening can be effective, however, there is a lack of evidence showing the superiority of any one type of exercise.^{74–76} Eccentric exercise was found equally effective as a conventional exercise programme for the rotator cuff and scapular muscles over 12

weeks in one RCT.⁷⁷ However, ESWT has emerged as an alternative efficacious and safe treatment option prior to considering surgery.⁷⁸ Several systematic reviews have concluded that ESWT is safe and effective in calcific rotator cuff tendinopathy, with ESWT improving shoulder function, reducing pain, and disintegrating calcific deposits, with efficacy maintained at 6-month follow-up.^{79,80} Despite compelling evidence and recommendations for calcific rotator cuff tendinopathy, none of these reviews or others recommend ESWT for non-calcific rotator cuff tendinopathy.⁶⁴ Supervised exercise has been found equally effective as ESWT for rotator cuff tendinopathy, therefore combining ESWT and exercise in a comprehensive rehabilitation program may lead to superior outcomes.⁸¹

Three studies have investigated ESWT combined with exercise for rotator cuff tendinopathy, with encouraging results. Kvalvaag et al.⁸² compared R-ESWT and supervised exercise for 12-weeks with sham R-ESWT and exercise in an RCT with 142 patients. At 24 and 52 weeks, participants in both groups had improved shoulder pain and function, but there were no differences between the groups. Prespecified subgroup analysis of patients with calcification in rotator cuff showed that the R-ESWT plus exercise group had a greater improvement in pain and function after 24 weeks but no difference was found at 52 weeks. Santamato et al.⁸³ compared three treatment sessions of F-ESWT with the same protocol plus ten supervised sessions of isokinetic exercise in an RCT with 30 patients. At two months follow-up, participants in the F-ESWT plus exercise group showed significantly less pain and greater improvement in function and muscle endurance than the F-ESWT group. The combined group was considered superior in the short to medium term, compared with F-ESWT alone. Carlisi et al.⁸⁴ compared F-ESWT plus a supervised eccentric training of the shoulder abductor muscles with F-ESWT only in 22 patients. At 9-weeks follow-up, there was a significant decrease in pain and an improvement in upper limb function in both groups, but no statistically significant differences between groups. However, as a pilot study with a small sample size, limited conclusions can be drawn.

Lateral elbow tendinopathy

Lateral elbow tendinopathy, commonly known as ‘tennis elbow’, arises from overuse and repetitive microtrauma of the extensor carpi radialis brevis muscle and tendon.⁸⁵ The term ‘Lateral epicondylitis’ refers to an inflammatory condition, however lateral elbow tendinopathy refers to chronic tendon degeneration.⁸⁶ Symptoms include lateral elbow pain which can radiate down the arm and weak grip strength.⁸⁷ Recreational sports, increased age, smoking, diabetes, repetitive work tasks using the hands or wrists are risk factors.⁸⁸ Annual incidence is 1%–3% and symptoms typically last 9–12 months, with most cases responding well to targeted exercise and few requiring surgeries.⁸⁹ Eccentric exercise has been shown to be effective for improving pain and function and is considered the gold standard conservative treatment.⁹⁰ However, exercise applied as monotherapy is considered to have lower effectiveness than when combined with other treatments such as ESWT.⁹¹ Although ESWT has been shown to be an efficacious treatment for lateral elbow tendinopathy, it is not without controversy as some studies have provided conflicting results.⁹² Despite methodological heterogeneity in many studies, systematic reviews have concluded there is the effectiveness of ESWT for lateral elbow tendinopathy, when studies have well-defined restrictive study parameters.⁹³

Three studies have investigated ESWT combined with exercise for lateral elbow tendinopathy, with all finding positive results. Celik et al.⁹⁴ compared photobiomodulation therapy applied three times a week for four weeks with F-ESWT once a week for four weeks in an RCT with 43 patients. Stretching and eccentric strengthening exercises were also given to both groups as a home program. Improvements for elbow extension and shoulder flexion strength and for painful movement were observed only in the photobiomodulation therapy group, whereas improvement of handgrip strength was present in both groups. Five

patients in the ESWT group reported no change, and 18 patients reported improvement compared with pre-treatment status. Both treatments were recommended, despite photobiomodulation therapy having better outcomes. Eraslan et al.⁹⁵ conducted an RCT with 45 patients randomised into three groups. Group one received 15 physiotherapy sessions including a home exercise programme of stretching and eccentric strengthening exercises. Group two received physiotherapy plus kinesio taping and group three received physiotherapy plus ESWT applied 3 times for 3 weeks. Pain intensity decreased, whereas maximum grip strength and functionality increased in all groups at the end of the treatment. The kinesio taping group and ESWT group yielded better results in improving function than the physiotherapy only group. A case series including 60 patients found that four sessions of F-ESWT combined with eccentric training four times a week for one month was effective for pain and function improvement at one, six, and 12 months after treatment.⁹⁶

Patellar tendinopathy

The aetiology of patellar tendinopathy or ‘jumpers’ knee’ involves extrinsic and intrinsic factors, with the pathogenesis-related to repetitive microtrauma to the patella tendon leading to an initial inflammatory response progressing to chronic degeneration.⁵⁹ Patellar tendinopathy is common in sports such as soccer, basketball, and volleyball, where prevalence has been found to be as high as 30%–45% in elite athletes.⁹⁷ Prevalence in the general population is thought to be around 14%.⁹⁸ Despite being challenging to treat, heavy slow resistance training and eccentric loading exercise can be effective, particularly eccentric decline squats.^{99,100} Wang et al.¹⁰¹ found that ESWT was more effective than stretching and strengthening exercises for the quadriceps and hamstrings. However, the authors did not give details of the type of exercises, so it is unclear if the recommended eccentric training was performed, which limits conclusions. Although studies have found ESWT to have positive results, there are controversies regarding treatment protocols, with published studies using extremely variable parameters.¹⁰² A systematic review including seven studies on ESWT for patellar tendinopathy, concluded that ESWT is an effective short and long-term treatment.²⁶ However, another systematic review concluded there was low-level evidence that ESWT is superior to standard conservative treatment in long-term outcomes.²⁷ Despite conflicting findings, all RCTs with robust methodological design have found positive effects of ESWT, recommending its use for patellar tendinopathy.⁹⁸

Four studies have investigated combined ESWT and exercise interventions for patellar tendinopathy, with the majority finding positive outcomes but not finding an additional benefit of ESWT over eccentric exercise. van Rijn et al.¹⁰³ conducted a secondary analysis of the combined databases of three double-blind RCTs, totalling 138 patients. Participants were divided into five groups based on treatment: ESWT, ESWT plus eccentric training, eccentric training, topical glyceryl trinitrate patch plus eccentric training, and placebo treatment. Fifty-two patients (38%) improved clinically after three months of treatment, with clinical improvement significantly higher in the eccentric training group and the ESWT plus eccentric training group compared with the other groups. Eccentric training was considered the most effective treatment. Thijs et al.¹⁰⁴ conducted an RCT with 52 patients, randomised to either F-ESWT or sham F-ESWT applied in three sessions at 1-week intervals and all participants were instructed to perform eccentric decline squat exercises daily for three months. No significant differences were found between the groups at six, 12, and 24 weeks, and no additional effect on ESWT to eccentric exercises was found. However, the results should be interpreted with caution due to a small sample size and considerable loss to follow-up, particularly in the ESWT group ($n = 22$).

Van der Worp et al.¹⁰⁵ conducted an RCT with 43 patients comparing three sessions of F-ESWT with three sessions of R-ESWT, with both groups receiving an eccentric training programme based on single-leg decline squats. Both groups improved significantly at 14 weeks

follow-up, but there was no statistically significant difference between groups. This was also the case for pain during sports activities and the decline squat. The results suggest that the type of ESWT may not be as important a consideration in patellar tendinopathy treatment as the inclusion of eccentric training. Vetrano et al.¹⁰⁶ conducted an RCT with 46 basketball, volleyball, and soccer athletes comparing two platelet-rich plasma injections over two weeks, with three sessions of F-ESWT. Both groups received a standardized stretching and strengthening exercise protocol. However, this protocol was only followed for two weeks, whereas recommended patellar tendinopathy strengthening programs are 12 weeks in duration.¹⁰⁰ Patients in both groups showed statistically significant improvement of symptoms at two, six, and 12-month follow-up, with no significant differences between groups at two-month follow-up. However, the platelet-rich plasma group showed significantly greater improvement at six and 12-month follow-up compared to the ESWT group in all outcome measures.

Achilles tendinopathy

Achilles tendinopathy is caused by overuse and repetitive microtrauma of the Achilles tendon and failed healing, leading to chronic degeneration.¹⁰⁷ The term Achilles ‘tendinitis’ was previously used to describe an acute local inflammatory response, which is histologically absent in chronic Achilles tendinopathy.¹⁰⁸ Clinical risk factors include impaired biomechanics, overtraining, high body mass index and fluoroquinolone use.¹⁰⁹ Chronic diseases and inflammatory conditions are also implicated, responsible for up to 2%–30% of cases.¹¹⁰ Symptoms include pain after activity or on palpation, swelling, and stiffness after rest, with up to 15% of patients being symptomatic five years after diagnosis.¹¹¹ Specific eccentric exercise protocols and heavy slow resistance training protocols have been shown to be effective and are widely recommended.¹¹² In unresponsive cases surgery may be required, however, ESWT has emerged as a potential alternative treatment.¹¹³ Eccentric exercise is considered the gold standard treatment for Achilles tendinopathy, although there is moderate evidence that ESWT may be more effective for insertional Achilles tendinopathy and equally effective for midportion Achilles tendinopathy.²⁷ Systematic reviews have found that ESWT is an effective short-term intervention for pain and function in Achilles tendinopathy, with low-quality evidence for long-term effectiveness.²⁶ Another systematic review concluded there was moderate evidence for F-ESWT in midportion and insertional Achilles tendinopathy.⁶⁴

Five studies have investigated combined ESWT and exercise interventions for Achilles tendinopathy, with all reporting superior outcomes for pain and function with the combined approach. An RCT with 68 patients by Rompe et al.¹¹⁴ compared a 12-week eccentric exercise protocol with the same protocol plus R-ESWT for three sessions. At four months from baseline, both groups had significantly improved pain and function, with 19 of 34 patients in group one (56%) and 28 of 34 patients in group two (82%) reporting being completely recovered or much improved. For all outcome measures, groups differed significantly in favour of the combined approach at the four-month follow-up. However, one year from baseline, there was no significant difference between groups. A prospective cohort study by Mansur et al.¹¹⁵ with 19 patients investigated a 12-week eccentric strengthening protocol combined with two sessions of R-ESWT. Fifteen (79%) patients were fully adherent to the exercise protocol, and 13 (68%) patients considered the treatment protocol successful. At 24 weeks follow-up, patients had significantly less pain and improved function, with the combined treatment being considered effective. Wheeler et al.¹¹⁶ conducted a prospective cohort of 63 patients receiving either three sessions of ESWT or a single high-volume image-guided injection. Both groups received a home exercise program based on eccentric strengthening. Statistically significant improvements in pain and function were found in both groups at three months follow-up, with no statistically significant differences between groups.

A case series by Wheeler et al.¹¹⁷ included 39 patients receiving R-ESWT once per week for three weeks, alongside a home exercise program including stretching, isometric and eccentric strengthening, core stability, and proprioception exercises. Statistically significant improvements were seen in insertional Achilles tendinopathy across a range of outcome measures for pain and function, however, these were less apparent for non-insertional tendinopathy. Pavone et al.¹¹⁸ conducted a case series with 40 patients who were previously unsuccessfully treated with eccentric exercises for three months. Patients underwent four sessions of F-ESWT with a two-week interval, together with eccentric exercises. At 12-month follow-up, 26 (65%) patients did not complain about pain, 11 (28%) patients got back to normal activities despite residual pain, and three (8%) patients still complained about pain. There had been no significant improvement in patients after previously completing eccentric exercises alone, with the combined ESWT and eccentric exercise intervention being recommended.

Plantar heel pain

Plantar heel pain (PHP) is one of the most prevalent musculoskeletal disorders, affecting up to 10% of the population, and causing 15% of all clinical foot symptoms.¹¹⁹ Degenerative shortening of the plantar fascia collagen matrix due to repetitive microtrauma is considered the pathophysiological origin of the condition, which can lead to heel spur formation.¹²⁰ Symptoms of PHP include morning heel pain and functional limitations such as impaired gait and reduced physical activities.¹²¹ The incidence of PHP in running athletes ranges from 5% to 10% and the prevalence ranges from 5% to 18%.¹²² Prior to the adoption of the clinical term PHP, the condition was previously referred to as ‘plantar fasciitis’ suggesting an inflammatory cause, and although inflammatory cells are present, degeneration due to repetitive microtrauma is considered the primary cause of the condition.¹²³ Moreover, the term ‘plantar fasciopathy’ is considered a more accurate diagnosis describing the chronic degenerative changes of the fascia, with many experts considering the condition to present like a tendinopathy.¹²⁰

For many years plantar fascia specific stretching (PFSS) has been the recommended exercise intervention for PHP due to being found effective in earlier RCTs.¹²⁴ However, heavy slow resistance training targeting the plantar fascia has been found more effective compared to PFSS for pain and function.³⁰ A systematic review concluded there is limited evidence for PFSS in isolation and combining other treatments with PFSS such as ESWT is recommended to increase effectiveness.¹²⁵ A methodologically robust meta-analysis by Sun et al.¹²⁶ included nine high-quality RCTs on ESWT and PHP. The review concluded that ESWT is associated with higher success rates and pain reduction compared to placebo in PHP, recommending it when conservative treatment fails. A recent network meta-analysis compared pain relief performance of eight different PHP therapies and included 41 studies and 2880 patients. In terms of one-month, three-month, and six-month pain outcomes, only ESWT provided better efficacy than placebo and ranked first for all seven outcomes and was recommended as the optimal treatment.¹²⁷

Ten RCTs, and one case series have investigated the effects of combined ESWT and exercise in PHP. All studies have used PFSS as an exercise intervention, with only the case series also using strengthening exercise. Rompe et al.¹²⁸ found that R-ESWT combined with PFSS was more effective than R-ESWT alone for pain in PHP at two and four months, but there was no difference at 24 months. Cinar et al.⁶⁸ compared the effectiveness of ESWT, low-level-laser therapy (LLLT), and control, with each group receiving home PFSS and insoles. There was a significant improvement in pain at three-months in all groups, both ESWT and LLLT were superior to control, with LLLT being superior to ESWT. Grecco et al.¹²⁹ found that R-ESWT combined with PFSS was superior for short term pain and function compared to ultrasound and PFSS, however, there was no difference at one-year. A three-arm RCT by Chew et al.¹³⁰ found that autologous blood injection with PFSS and ESWT with PFSS were both more effective than PFSS alone for pain and

function in PHP at six-months.

A three-arm RCT by Ulusoy et al.¹³¹ compared ESWT, ultrasound and LLLT, all combined with PFSS. All three groups significantly improved PHP pain and function at one-month, with treatment success determined as 71% for LLLT, 65% for ESWT and 24% for the ultrasound. Okur et al.¹³² compared R-ESWT and daily PFSS with custom-fabricated orthotics and PFSS over four weeks. Both groups achieved significant improvements in PHP pain at 24 weeks, with no significant difference between the groups. Eslamian et al.¹³³ compared ESWT and PFSS with a single corticosteroid injection and PFSS, finding significant PHP pain and function improvement in both groups, although the ESWT group had better outcomes (67% versus 48%). Good or excellent results in the opinions of patients were achieved in 55% of ESWT and 30% of corticosteroid injection groups. A three-arm RCT by Akinoglu et al.¹³⁴ compared R-ESWT combined with PFSS with ultrasound and PFSS and PFSS alone. At one-month all groups had improved PHP pain and function, with both the R-ESWT and ultrasound groups being superior to PFSS alone, and ankle proprioception sense increased only in the R-ESWT group. Vahdatpour et al.¹³⁵ compared ESWT combined with PFSS and topical corticosteroid versus ESWT combined with PFSS and sham corticosteroid. Both groups improved pain and function at one and three months, with the topical corticosteroid group being superior at one month, with no difference between groups at three months. The study identified how topical corticosteroids could enhance the short-term effectiveness of combined ESWT and PFSS.

A four-arm RCT by Takla et al.¹³⁶ compared ESWT combined with PFSS and LLLT, with each of ESWT and LLLT combined with PFSS, and sham LLLT with PFSS. All groups except the sham LLLT group showed significant pain and function improvement at three-months, with the combined ESWT, PFSS, and LLLT group being most effective. Combined ESWT and PFSS was found to be superior to LLLT and sham LLLT combined with PFSS. The study was the first to show that combining LLLT with ESWT and PFSS can lead to improved outcomes in PHP. A recent case series found that ESWT plus a progressive home exercise program including PFSS and strength exercise, led to significantly improved function and pain but not increased activity levels.¹³⁷ Specific details of the strength exercise were omitted and despite being a case series with no comparator and only 35 participants, the study demonstrated the feasibility of combined ESWT and multiple types of exercise. A recent RCT found that combining heavy slow resistance training and PFSS with corticosteroid injection, was significantly more effective than corticosteroid injection or the combined exercise program alone for pain and function in PHP.¹³⁸ However, no RCTs have investigated the effects of a combined ESWT and heavy slow resistance training intervention for PHP.¹³⁹

Conclusion

Tendinopathy has a high prevalence in the general population and in athletes, with both exercise and ESWT in isolation being found to be effective treatments in several studies. A limited number of studies have investigated combined exercise and ESWT for common tendinopathies, with further large high-quality RCTs required. The currently limited evidence for combined ESWT and exercise interventions is positive for PHP, Achilles, lateral elbow, and rotator cuff tendinopathy, especially when calcification exists. Despite overall positive outcomes in patellar tendinopathy, the combined treatment has not been found to offer additional benefits over eccentric training alone. However, studies not showing additional benefits have had methodological limitations and small sample sizes, limiting conclusions. Current evidence recommends combined rather than single modalities in tendinopathy treatment to achieve superior long-term outcomes. However, there is a dearth of high-quality RCTs investigating combined interventions for tendinopathies, such as ESWT, exercise, and other emerging treatments. A clear need exists for further studies comparing combined treatments, such as comprehensive exercise programs which include different types of

exercise as opposed to only one type. Detailed description of exercise protocols, adherence, and progression parameters are required in future studies, with a lack of information provided in current studies. The current encouraging evidence suggests that combined specific exercise and ESWT interventions should be recommended for PHP, rotator cuff, lateral elbow, and Achilles tendinopathies. Further well-designed RCTs are required to make definitive recommendations on the optimal treatment protocols for tendinopathies, particularly patellar tendinopathy.

Key recommendations

- Eccentric exercise combined with ESWT is feasible and may be more effective than either alone for Achilles and lateral elbow tendinopathies and may therefore be recommended as combined treatment.
- Isotonic or eccentric exercise combined with ESWT is more effective than either alone for rotator cuff tendinopathy, particularly when calcification exists, and may therefore be recommended.
- ESWT does not provide any additional effect to eccentric exercise for patellar tendinopathy, therefore either treatment in isolation is recommended, with eccentric exercise the most recommended treatment.
- ESWT combined with PFSS can be recommended for treating PHP as it is more effective than either alone. The effectiveness of the combination can be increased further by combining it with LLLT or topical corticosteroid. The effects of combined ESWT and resistance training have not been investigated.
- Overall, tendon specific exercise should be the recommended treatment for tendinopathy in isolation, with combined exercise and ESWT offered when available. Although ESWT can be effective in isolation, combining it with specific exercise will likely be more effective. ESWT should not be recommended over tendon-specific exercise unless exercise is not appropriate for individual patients.

Submission statement

The work described has not been published previously, it is not under consideration for publication elsewhere, its publication is approved by all authors, and that, if accepted, it will not be published elsewhere including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Conflict of interest

The author declares no conflicts of interest relevant to the content of this review.

References

1. Docking SI, Cook J. How do tendons adapt? Going beyond tissue responses to understand positive adaptation and pathology development: a narrative review. *J Musculoskelet Neuronal Interact.* 2019;19(3):300–310.
2. Singh A, Calafi A, Diefenbach C, Kreulen C, Giza E. Noninsertional tendinopathy of the achilles. *Foot Ankle Clin.* 2017;22(4):745–760. <https://doi.org/10.1016/j.fcl.2017.07.006>.
3. Magnusson SP, Langberg H, Kjaer M. The pathogenesis of tendinopathy: balancing the response to loading. *Nat Rev Rheumatol.* 2010;6(5):262–268. <https://doi.org/10.1038/nrrheum.2010.43>.
4. Malliaras P, Barton CJ, Reeves ND, Langberg H. Achilles and patellar tendinopathy loading programmes : a systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sports Med.* 2013;43(4): 267–286. <https://doi.org/10.1007/s40279-013-0019-z>.
5. Fu SC, Rolf C, Cheuk YC, Lui PP, Chan KM. Deciphering the pathogenesis of tendinopathy: a three-stages process. *Sports Med Arthrosc Rehabil Ther Technol.* 2010;2:30. <https://doi.org/10.1186/1758-2555-2-30>.
6. Rees JD, Stride M, Scott A. Tendons—time to revisit inflammation. *Br J Sports Med.* 2014;48(21):1553–1557. <https://doi.org/10.1136/bjsports-2012-091957>.
7. Riel H, Lindstrøm CF, Rathleff MS, Jensen MB, Olesen JL. Prevalence and incidence rate of lower-extremity tendinopathies in a Danish general practice: a registry-based study. *BMC Musculoskel Disord.* 2019;20(1):239. <https://doi.org/10.1186/s12891-019-2629-6>.

8. Albers IS, Zwerver J, Dierckx RL, Dekker JH, Van den Akker-Scheek I. Incidence and prevalence of lower extremity tendinopathy in a Dutch general practice population: a cross sectional study. *BMC Musculoskel Disord.* 2016;17:16. <https://doi.org/10.1186/s12891-016-0885-2>.
9. Hopkins C, Fu SC, Chua E, et al. Critical review on the socio-economic impact of tendinopathy. *Asia Pac J Sports Med Arthrosc Rehabil Technol.* 2016;4:9–20. <https://doi.org/10.1016/j.asmart.2016.01.002>.
10. Monteleone G, Tramontana A, Mc Donald K, Sorge R, Tiloca A, Foti C. Ultrasonographic evaluation of the shoulder in elite Italian beach volleyball players. *J Sports Med Phys Fit.* 2015;55(10):1193–1199.
11. Pieber K, Angelmaier L, Csapo R, Herczeg M. Acute injuries and overuse syndromes in sport climbing and bouldering in Austria: a descriptive epidemiological study. *Wien Klin Wochenschr.* 2012;124(11-12):357–362. <https://doi.org/10.1007/s00508-012-0174-5>.
12. McCarthy MM, Voos JE, Nguyen JT, Callahan L, Hannafin JA. Injury profile in elite female basketball athletes at the Women's National Basketball Association combine. *Am J Sports Med.* 2013;41(3):645–651. <https://doi.org/10.1177/0363546512474223>.
13. Cassel M, Baur H, Hirschmüller A, Carlsohn A, Fröhlich K, Mayer F. Prevalence of Achilles and patellar tendinopathy and their association to intratendinous changes in adolescent athletes. *Scand J Med Sci Sports.* 2015;25(3):e310–e318. <https://doi.org/10.1111/sms.12318>.
14. Buda R, Di Caprio F, Bedetti L, Mosca M, Giannini S. Foot overuse diseases in rock climbing: an epidemiologic study. *J Am Podiatr Med Assoc.* 2013;103(2):113–120. <https://doi.org/10.7547/1030113>.
15. Tenforde AS, Sayres LC, McCurdy ML, Collado H, Sainani KL, Fredericson M. Overuse injuries in high school runners: lifetime prevalence and prevention strategies. *Pharm Manag PM R.* 2011;3(2):125–131. <https://doi.org/10.1016/j.pmrj.2010.09.009>.
16. Seitz AL, McClure PW, Finucane S, Boardman 3rd ND, Michener LA. Mechanisms of rotator cuff tendinopathy: intrinsic, extrinsic, or both? *Clin Biomech.* 2011;26(1):1–12. <https://doi.org/10.1016/j.clinbiomech.2010.08.001>.
17. Wheeler PC. Up to a quarter of patients with certain chronic recalcitrant tendinopathies may have central sensitisation: a prospective cohort of more than 300 patients. *Br J Pain.* 2019;13(3):137–144. <https://doi.org/10.1177/2049463718800352>.
18. Plinsinga ML, Coombes BK, Mellor R, et al. Psychological factors not strength deficits are associated with severity of gluteal tendinopathy: a cross-sectional study. *Eur J Pain.* 2018;22(6):1124–1133. <https://doi.org/10.1002/ejp.1199>.
19. Plinsinga ML, Brink MS, Vicenzino B, van Wilgen CP. Evidence of nervous system sensitization in commonly presenting and persistent painful tendinopathies: a systematic review. *J Orthop Sports Phys Ther.* 2015;45(11):864–875. <https://doi.org/10.2519/jospt.2015.5895>.
20. Sprague AL, Smith AH, Knox P, Pohlig RT, Grävare Silbernagel K. Modifiable risk factors for patellar tendinopathy in athletes: a systematic review and meta-analysis. *Br J Sports Med.* 2018;52(24):1575–1585. <https://doi.org/10.1136/bjsports-2017-099000>.
21. Abate M, Schiavone C, Salini V, Andia I. Occurrence of tendon pathologies in metabolic disorders. *Rheumatology.* 2013;52(4):599–608. <https://doi.org/10.1093/rheumatology/kes395>.
22. Scott A, Zwerver J, Grewal N, et al. Lipids, adiposity and tendinopathy: is there a mechanistic link? Critical review. *Br J Sports Med.* 2015;49(15):984–988. <https://doi.org/10.1136/bjsports-2014-093989>.
23. Alves C, Mendes D, Marques FB. Fluoroquinolones and the risk of tendon injury: a systematic review and meta-analysis. *Eur J Clin Pharmacol.* 2019;75(10):1431–1443. <https://doi.org/10.1007/s00228-019-02713-1>.
24. Cardoso TB, Pizzari T, Kinsella R, Hope D, Cook JL. Current trends in tendinopathy management. *Best Pract Res Clin Rheumatol.* 2019;33(1):122–140. <https://doi.org/10.1016/j.berh.2019.02.001>.
25. Dimitrios S. Exercise for tendinopathy. *World J Methodol.* 2015;5(2):51–54. <https://doi.org/10.5662/wjm.v5.i2.51>.
26. Mani-Babu S, Morrissey D, Waugh C, Screen H, Barton C. The effectiveness of extracorporeal shock wave therapy in lower limb tendinopathy: a systematic review. *Am J Sports Med.* 2015;43(3):752–761. <https://doi.org/10.1177/0363546514531911>.
27. Korakakis V, Whiteley R, Tzavara A, Malliaropoulos N. The effectiveness of extracorporeal shockwave therapy in common lower limb conditions: a systematic review including quantification of patient-rated pain reduction. *Br J Sports Med.* 2018;52(6):387–407. <https://doi.org/10.1136/bjsports-2016-097347>.
28. Romeo P, Lavanga V, Pagani D, Sansone V. Extracorporeal shock wave therapy in musculoskeletal disorders: a review. *Med Princ Pract.* 2014;23(1):7–13. <https://doi.org/10.1159/000355472>.
29. Wang CJ. Extracorporeal shockwave therapy in musculoskeletal disorders. *J Orthop Surg Res.* 2012;7:11. <https://doi.org/10.1186/1749-799X-7-11>.
30. Rathleff MS, Mølgaard CM, Fredberg U, et al. High-load strength training improves outcome in patients with plantar fasciitis: a randomized controlled trial with 12-month follow-up. *Scand J Med Sci Sports.* 2015;25(3):e292–e300. <https://doi.org/10.1111/sms.12313>.
31. Holmgren T, Björnsson Hallgren H, Öberg B, Adolfsson L, Johansson K. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. *BMJ.* 2012;344:e787. <https://doi.org/10.1136/bmj.e787>.
32. Stasinopoulos D, Stasinopoulos I. Comparison of effects of eccentric training, eccentric-concentric training, and eccentric-concentric training combined with isometric contraction in the treatment of lateral elbow tendinopathy. *J Hand Ther.* 2017;30(1):13–19. <https://doi.org/10.1016/j.jht.2016.09.001>.
33. Beyer R, Kongsgaard M, Hougs Kjær B, Øhlenschläger T, Kjær M, Magnusson SP. Heavy slow resistance versus eccentric training as treatment for Achilles tendinopathy: a randomized controlled trial. *Am J Sports Med.* 2015;43(7):1704–1711. <https://doi.org/10.1177/0363546515584760>.
34. Kongsgaard M, Kovanen V, Aagaard P, et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports.* 2009;19(6):790–802. <https://doi.org/10.1111/j.1600-0838.2009.00949.x>.
35. Kongsgaard M, Qvortrup K, Larsen J, et al. Fibril morphology and tendon mechanical properties in patellar tendinopathy: effects of heavy slow resistance training. *Am J Sports Med.* 2010;38(4):749–756. <https://doi.org/10.1177/0363546509350915>.
36. Murtaugh B, Ihm JM. Eccentric training for the treatment of tendinopathies. *Curr Sports Med Rep.* 2013;12(3):175–182. <https://doi.org/10.1249/JSR.0b013e3182933761>.
37. Khan KM, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med.* 2009;43(4):247–252. <https://doi.org/10.1136/bjism.2008.054239>.
38. McAuliffe S, Tabuena A, McCreesh K, et al. Altered strength profile in achilles tendinopathy: a systematic review and meta-analysis. *J Athl Train.* 2019;54(8):889–900. <https://doi.org/10.4085/1062-6050-43-18>.
39. Wang JH, Guo Q, Li B. Tendon biomechanics and mechanobiology—a minireview of basic concepts and recent advancements. *J Hand Ther.* 2012;25(2):133–141. <https://doi.org/10.1016/j.jht.2011.07.004>.
40. Wearing SC, Grigg NL, Hooper SL, Smeathers JE. Conditioning of the Achilles tendon via ankle exercise improves correlations between sonographic measures of tendon thickness and body anthropometry. *J Appl Physiol (1985).* 2011;110(5):1384–1389. <https://doi.org/10.1152/japplphysiol.00075.2011>.
41. Drew BT, Smith TO, Littlewood C, Sturrock B. Do structural changes (eg, collagen/matrix) explain the response to therapeutic exercises in tendinopathy: a systematic review. *Br J Sports Med.* 2014;48(12):966–972. <https://doi.org/10.1136/bjsports-2012-091285>.
42. Heinemeier KM, Schjerling P, Heinemeier J, Magnusson SP, Kjær M. Lack of tissue renewal in human adult Achilles tendon is revealed by nuclear bomb (14)C. *FASEB J.* 2013;27(5):2074–2079. <https://doi.org/10.1096/fj.12-225599>.
43. Kongsgaard M, Reitelsheder S, Pedersen TG, et al. Region specific patellar tendon hypertrophy in humans following resistance training. *Acta Physiol.* 2007;191(2):111–121. <https://doi.org/10.1111/j.1748-1716.2007.01714.x>.
44. Bohm S, Mersmann F, Arampatzis A. Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults. *Sports Med Open.* 2015;1(1):7. <https://doi.org/10.1186/s40798-015-0009-9>.
45. Ioppolo F, Rompe JD, Furia JP, Cacchio A. Clinical application of shock wave therapy (SWT) in musculoskeletal disorders. *Eur J Phys Rehabil Med.* 2014;50(2):217–230.
46. Liao CD, Tsauo JY, Chen HC, Liou TH. Efficacy of extracorporeal shock wave therapy for lower-limb tendinopathy: a meta-analysis of randomized controlled trials. *Am J Phys Med Rehabil.* 2018;97(9):605–619. <https://doi.org/10.1097/PHM.0000000000000925>.
47. Testa G, Vescio A, Perez S, et al. Extracorporeal shockwave therapy treatment in upper limb diseases: a systematic review. *J Clin Med.* 2020;9(2):453. <https://doi.org/10.3390/jcm9020453>.
48. d'Agostino MC, Craig K, Tibalt E, Respizzi S. Shock wave as biological therapeutic tool: from mechanical stimulation to recovery and healing, through mechanotransduction. *Int J Surg.* 2015;24(Pt B):147–153. <https://doi.org/10.1016/j.ijsu.2015.11.030>.
49. Foldager CB, Kearney C, Spector M. Clinical application of extracorporeal shock wave therapy in orthopedics: focused versus unfocused shock waves. *Ultrasound Med Biol.* 2012;38(10):1673–1680. <https://doi.org/10.1016/j.ultrasmedbio.2012.06.004>.
50. Moya D, Ramón S, Schaden W, Wang CJ, Guiloff L, Cheng JH. The role of extracorporeal shockwave treatment in musculoskeletal disorders. *J Bone Joint Surg Am.* 2018;100(3):251–263. <https://doi.org/10.2106/JBJS.17.00661>.
51. Wang CJ, Cheng JH, Chou WY, Hsu SL, Chen JH, Huang CY. Changes of articular cartilage and subchondral bone after extracorporeal shockwave therapy in osteoarthritis of the knee. *Int J Med Sci.* 2017;14(3):213–223. <https://doi.org/10.7150/ijms.17469>.
52. Holfeld J, Tepeköylü C, Blunder S, et al. Low energy shock wave therapy induces angiogenesis in acute hind-limb ischemia via VEGF receptor 2 phosphorylation. *PLoS One.* 2014;9(8):e103982. <https://doi.org/10.1371/journal.pone.0103982>.
53. Tara S, Miyamoto M, Takagi G, et al. Low-energy extracorporeal shock wave therapy improves microcirculation blood flow of ischemic limbs in patients with peripheral arterial disease: pilot study. *J Nippon Med Sch.* 2014;81(1):19–27. <https://doi.org/10.1272/jnms.81.19>.
54. Abe Y, Ito K, Hao K, et al. Extracorporeal low-energy shock-wave therapy exerts anti-inflammatory effects in a rat model of acute myocardial infarction. *Circ J.* 2014;78(12):2915–2925. <https://doi.org/10.1253/circj.14-0230>.
55. Chen YL, Chen KH, Yin TC, et al. Extracorporeal shock wave therapy effectively prevented diabetic neuropathy. *Am J Transl Res.* 2015;7(12):2543–2560.
56. Davis TA, Stojadinovic A, Anam K, et al. Extracorporeal shock wave therapy suppresses the early proinflammatory immune response to a severe cutaneous burn injury. *Int Wound J.* 2009;6(1):11–21. <https://doi.org/10.1111/j.1742-481X.2008.00540.x>.
57. Wess OJ. A neural model for chronic pain and pain relief by extracorporeal shock wave treatment. *Urol Res.* 2008;36(6):327–334. <https://doi.org/10.1007/s00240-008-0156-2>.

58. Waugh CM, Morrissey D, Jones E, Riley GP, Langberg H, Screen HR. In vivo biological response to extracorporeal shockwave therapy in human tendinopathy. *Eur Cell Mater.* 2015;29:268–280. <https://doi.org/10.22203/ecm.v029a20>.
59. Malliaras P, Cook J, Purdam C, Rio E. Patellar tendinopathy: clinical diagnosis, load management, and advice for challenging case presentations. *J Orthop Sports Phys Ther.* 2015;45(11):887–898. <https://doi.org/10.2519/jospt.2015.5987>.
60. Lohrer H, Nauck T, Korakakis V, Malliaropoulos N. Historical ESWT paradigms are overcome: a narrative review. *BioMed Res Int.* 2016;2016:3850461. <https://doi.org/10.1155/2016/3850461>.
61. Cook JL, Rio E, Purdam CR, Docking SI. Revisiting the continuum model of tendon pathology: what is its merit in clinical practice and research? *Br J Sports Med.* 2016;50(19):1187–1191. <https://doi.org/10.1136/bjsports-2015-095422>.
62. Cook JL, Purdam C. Is compressive load a factor in the development of tendinopathy? *Br J Sports Med.* 2012;46(3):163–168. <https://doi.org/10.1136/bjsports-2011-090414>.
63. Gaida JE, Cook J. Treatment options for patellar tendinopathy: critical review. *Curr Sports Med Rep.* 2011;10(5):255–270. <https://doi.org/10.1249/JSR.0b013e31822d4016>.
64. Speed C. A systematic review of shockwave therapies in soft tissue conditions: focusing on the evidence. *Br J Sports Med.* 2014;48(21):1538–1542. <https://doi.org/10.1136/bjsports-2012-091961>.
65. Leung R, Malliaropoulos N, Korakakis V, Padhiar N. What are patients' knowledge, expectation and experience of radial extracorporeal shockwave therapy for the treatment of their tendinopathies? A qualitative study. *J Foot Ankle Res.* 2018;11:11. <https://doi.org/10.1186/s13047-018-0254-5>.
66. Sutton DA, Nordin M, Côté P, et al. The effectiveness of multimodal care for soft tissue injuries of the lower extremity: a systematic review by the Ontario Protocol for Traumatic Injury Management (OPTIMA) Collaboration. *J Manip Physiol Ther.* 2016;39(2):95–109. <https://doi.org/10.1016/j.jmpt.2016.01.004>.
67. Riel H, Jensen MB, Olesen JL, Vicenzino B, Rathleff MS. Self-dosed and pre-determined progressive heavy-slow resistance training have similar effects in people with plantar fasciopathy: a randomised trial. *J Physiother.* 2019;65(3):144–151. <https://doi.org/10.1016/j.jphys.2019.05.011>.
68. Cinar E, Saxena S, Uygur F. Combination therapy versus exercise and orthotic support in the management of pain in plantar fasciitis: a randomized controlled trial. *Foot Ankle Int.* 2018;39(4):406–414. <https://doi.org/10.1177/1071100717747590>.
69. Biberthaler P, Wiedemann E, Nerlich A, et al. Microcirculation associated with degenerative rotator cuff lesions. In vivo assessment with orthogonal polarization spectral imaging during arthroscopy of the shoulder. *J Bone Joint Surg Am.* 2003;85(3):475–480.
70. Kaux JF, Forthomme B, Goff CL, Crielaard JM, Croisier JL. Current opinions on tendinopathy. *J Sports Sci Med.* 2011;10(2):238–253.
71. de Witte PB, van Adrichem RA, Selten JW, et al. Persistent shoulder symptoms in calcific tendinitis: clinical and radiological predictors. *Ned Tijdschr Geneesk.* 2016;160:D521. <https://doi.org/10.1007/s00330-016-4224-7>.
72. Sansone V, Maiorano E, Galluzzo A, Pascale V. Calcific tendinopathy of the shoulder: clinical perspectives into the mechanisms, pathogenesis, and treatment. *Orthop Res Rev.* 2018;10:63–72. <https://doi.org/10.2147/ORR.S138225>.
73. Rebuzzi E, Coletti N, Schiavetti S, Giusto F. Arthroscopic surgery versus shock wave therapy for chronic calcifying tendinitis of the shoulder. *J Orthop Traumatol.* 2008;9(4):179–185. <https://doi.org/10.1007/s10195-008-0024-4>.
74. Shire AR, Stæhr TAB, Overby JB, Bastholm Dahl M, Sandell Jacobsen J, Høytrup Christiansen D. Specific or general exercise strategy for subacromial impingement syndrome—does it matter? A systematic literature review and meta-analysis. *BMC Musculoskel Disord.* 2017;18(1):158. <https://doi.org/10.1186/s12891-017-1518-0>. Published 2017 Apr 17.
75. Larsson R, Bernhardtsson S, Nordeman L. Effects of eccentric exercise in patients with subacromial impingement syndrome: a systematic review and meta-analysis. *BMC Musculoskel Disord.* 2019;20(1):446. <https://doi.org/10.1186/s12891-019-2796-5>.
76. Ingwersen KG, Jensen SL, Sørensen L, et al. Three months of progressive high-load versus traditional low-load strength training among patients with rotator cuff tendinopathy: primary results from the double-blind randomized controlled RoCTEx Trial. *Orthop J Sports Med.* 2017;5(8):2325967117723292. <https://doi.org/10.1177/2325967117723292>.
77. DeJaco B, Habets B, van Loon C, van Grinsven S, van Cingel R. Eccentric versus conventional exercise therapy in patients with rotator cuff tendinopathy: a randomized, single blinded, clinical trial. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(7):2051–2059. <https://doi.org/10.1007/s00167-016-4223-x>.
78. Louwerens JK, Veltman ES, van Noort A, van den Bekerom MP. The effectiveness of high-energy extracorporeal shockwave therapy versus ultrasound-guided needling versus arthroscopic surgery in the management of chronic calcific rotator cuff tendinopathy: a systematic review. *Arthroscopy.* 2016;32(1):165–175. <https://doi.org/10.1016/j.arthro.2015.06.049>.
79. Louwerens JK, Siersevelt IN, van Noort A, van den Bekerom MP. Evidence for minimally invasive therapies in the management of chronic calcific tendinopathy of the rotator cuff: a systematic review and meta-analysis. *J Shoulder Elbow Surg.* 2014;23(8):1240–1249. <https://doi.org/10.1016/j.jse.2014.02.002>.
80. Ioppolo F, Tattoli M, Di Sante L, et al. Clinical improvement and resorption of calcifications in calcific tendinitis of the shoulder after shock wave therapy at 6 months' follow-up: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* 2013;94(9):1699–1706. <https://doi.org/10.1016/j.apmr.2013.01.030>.
81. Engebretsen K, Grotle M, Bautz-Holter E, Ekeberg OM, Juel NG, Brox JI. Supervised exercises compared with radial extracorporeal shock-wave therapy for subacromial shoulder pain: 1-year results of a single-blind randomized controlled trial. *Phys Ther.* 2011;91(1):37–47. <https://doi.org/10.2522/ptj.20090338>.
82. Kvalvaag E, Roe C, Engebretsen KB, et al. One year results of a randomized controlled trial on radial Extracorporeal Shock Wave Treatment, with predictors of pain, disability and return to work in patients with subacromial pain syndrome. *Eur J Phys Rehabil Med.* 2018;54(3):341–350. <https://doi.org/10.23736/S1973-9087.17.04748-7>.
83. Santamato A, Panza F, Notarnicola A, et al. Is extracorporeal shockwave therapy combined with isokinetic exercise more effective than extracorporeal shockwave therapy alone for subacromial impingement syndrome? a randomized clinical trial. *J Orthop Sports Phys Ther.* 2016;46(9):714–725. <https://doi.org/10.2519/jospt.2016.4629>.
84. Carlisi E, Lisi C, Dall'angelo A, et al. Focused extracorporeal shock wave therapy combined with supervised eccentric training for supraspinatus calcific tendinopathy. *Eur J Phys Rehabil Med.* 2018;54(1):41–47. <https://doi.org/10.23736/S1973-9087.16.04299-4>.
85. Bhabra G, Wang A, Ebert JR, Edwards P, Zheng M, Zheng MH. Lateral elbow tendinopathy: development of a pathophysiology-based treatment algorithm. *Orthop J Sports Med.* 2016;4(11):2325967116670635. <https://doi.org/10.1177/2325967116670635>.
86. Thiele S, Thiele R, Gerdemeyer L. Lateral epicondylitis: this is still a main indication for extracorporeal shockwave therapy. *Int J Surg.* 2015;24(Pt B):165–170. <https://doi.org/10.1016/j.ijssu.2015.09.034>.
87. De Smedt T, de Jong A, Van Leemput W, Lieven D, Van Glabbeek F. Lateral epicondylitis in tennis: update on aetiology, biomechanics and treatment. *Br J Sports Med.* 2007;41(11):816–819. <https://doi.org/10.1136/bjsm.2007.036723>.
88. Shiri R, Viikari-Juntura E, Varonen H, Heliövaara M. Prevalence and determinants of lateral and medial epicondylitis: a population study. *Am J Epidemiol.* 2006;164(11):1065–1074. <https://doi.org/10.1093/aje/kwj325>.
89. Malliaras P, Maffulli N, Garau G. Eccentric training programmes in the management of lateral elbow tendinopathy. *Disabil Rehabil.* 2008;30(20-22):1590–1596. <https://doi.org/10.1080/09638280701786195>.
90. Cullinane FL, Boocock MG, Trevelyan FC. Is eccentric exercise an effective treatment for lateral epicondylitis? A systematic review. *Clin Rehabil.* 2014;28(1):3–19. <https://doi.org/10.1177/0269215513491974>.
91. Dimitrios S. Lateral elbow tendinopathy: evidence of physiotherapy management. *World J Orthoped.* 2016;7(8):463–466. <https://doi.org/10.5312/wjo.v7.i8.463>.
92. Notarnicola A, Maccagnano G, Tafuri S, et al. Prognostic factors of extracorporeal shock wave therapy for tendinopathies. *Musculoskel Surg.* 2016;100(1):53–61. <https://doi.org/10.1007/s12306-015-0375-y>.
93. Rompe JD, Maffulli N. Repetitive shock wave therapy for lateral elbow tendinopathy (tennis elbow): a systematic and qualitative analysis. *Br Med Bull.* 2007;83:355–378. <https://doi.org/10.1093/bmb/ldm019>.
94. Celik D, Anaforoglu Kulunkoglu B. Photobiomodulation therapy versus extracorporeal shock wave therapy in the treatment of lateral epicondylitis. *Photobiomodul Photomed Laser Surg.* 2019;37(5):269–275. <https://doi.org/10.1089/photob.2018.4533>.
95. Eraslan L, Yuce D, Erbilici A, Baltaci G. Does Kinesiotaping improve pain and functionality in patients with newly diagnosed lateral epicondylitis? *Knee Surg Sports Traumatol Arthrosc.* 2018;26(3):938–945. <https://doi.org/10.1007/s00167-017-4691-7>.
96. Testa G, Vescio A, Perez S, et al. Functional outcome at short and middle term of the extracorporeal shockwave therapy treatment in lateral epicondylitis: a case-series study. *J Clin Med.* 2020;9(3):633. <https://doi.org/10.3390/jcm9030633>.
97. van der Worp H, van Ark M, Roerink S, Pepping GJ, van den Akker-Scheek I, Zwerver J. Risk factors for patellar tendinopathy: a systematic review of the literature. *Br J Sports Med.* 2011;45(5):446–452. <https://doi.org/10.1136/bjsm.2011.084079>.
98. Leal C, Ramon S, Furia J, Fernandez A, Romero L, Hernandez-Sierra L. Current concepts of shockwave therapy in chronic patellar tendinopathy. *Int J Surg.* 2015;24(Pt B):160–164. <https://doi.org/10.1016/j.ijssu.2015.09.066>.
99. Schwartz A, Watson JN, Hutchinson MR. *Patellar Tendinopathy.* *Sports Health.* 2015;7(5):415–420. <https://doi.org/10.1177/1941738114568775>.
100. Everhart JS, Cole D, Sojka JH, et al. Treatment options for patellar tendinopathy: a systematic review. *Arthroscopy.* 2017;33(4):861–872. <https://doi.org/10.1016/j.arthro.2016.11.007>.
101. Wang CJ, Ko JY, Chan YS, Weng LH, Hsu SL. Extracorporeal shockwave for chronic patellar tendinopathy. *Am J Sports Med.* 2007;35(6):972–978. <https://doi.org/10.1177/0363546506298109>.
102. van der Worp H, van den Akker-Scheek I, van Schie H, Zwerver J. ESWT for tendinopathy: technology and clinical implications. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(6):1451–1458. <https://doi.org/10.1007/s00167-012-2009-3>.
103. van Rijn D, van den Akker-Scheek I, Steunebrink M, Diercks RL, Zwerver J, van der Worp H. Comparison of the effect of 5 different treatment options for managing patellar tendinopathy: a secondary analysis. *Clin J Sport Med.* 2019;29(3):181–187. <https://doi.org/10.1097/JSM.0000000000000520>.
104. Thijs KM, Zwerver J, Backx FJ, et al. Effectiveness of shockwave treatment combined with eccentric training for patellar tendinopathy: a double-blinded randomized study. *Clin J Sport Med.* 2017;27(2):89–96. <https://doi.org/10.1097/JSM.0000000000000332>.
105. van der Worp H, Zwerver J, Hamstra M, van den Akker-Scheek I, Diercks RL. No difference in effectiveness between focused and radial shockwave therapy for treating patellar tendinopathy: a randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc.* 2014;22(9):2026–2032. <https://doi.org/10.1007/s00167-013-2522-z>.

106. Vetrano M, Castorina A, Vulpiani MC, Baldini R, Pavan A, Ferretti A. Platelet-rich plasma versus focused shock waves in the treatment of jumper's knee in athletes. *Am J Sports Med.* 2013;41(4):795–803. <https://doi.org/10.1177/0363546513475345>.
107. Magnan B, Bondi M, Pierantoni S, Samaila E. The pathogenesis of Achilles tendinopathy: a systematic review. *Foot Ankle Surg.* 2014;20(3):154–159. <https://doi.org/10.1016/j.fas.2014.02.010>.
108. Lopez RG, Jung HG. Achilles tendinosis: treatment options. *Clin Orthop Surg.* 2015; 7(1):1–7. <https://doi.org/10.4055/cios.2015.7.1.1>.
109. van der Vlist AC, Breda SJ, Oei EHG, Verhaar JAN, de Vos RJ. Clinical risk factors for Achilles tendinopathy: a systematic review. *Br J Sports Med.* 2019;53(21): 1352–1361. <https://doi.org/10.1136/bjsports-2018-099991>.
110. Silbernagel KG, Crossley KM. A proposed return-to-sport program for patients with midportion Achilles tendinopathy: rationale and implementation. *J Orthop Sports Phys Ther.* 2015;45(11):876–886. <https://doi.org/10.2519/jospt.2015.5885>.
111. Silbernagel KG, Brorsson A, Lundberg M. The majority of patients with Achilles tendinopathy recover fully when treated with exercise alone: a 5-year follow-up. *Am J Sports Med.* 2011;39(3):607–613. <https://doi.org/10.1177/0363546510384789>.
112. Murphy MC, Travers MJ, Chivers P, et al. Efficacy of heavy eccentric calf training for treating mid-portion Achilles tendinopathy: a systematic review and meta-analysis. *Br J Sports Med.* 2019;53(17):1070–1077. <https://doi.org/10.1136/bjsports-2018-099934>.
113. Gerdemeyer L, Mittermayr R, Fuerst M, et al. Current evidence of extracorporeal shock wave therapy in chronic Achilles tendinopathy. *Int J Surg.* 2015;24(Pt B): 154–159. <https://doi.org/10.1016/j.ijsu.2015.07.718>.
114. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion Achilles tendinopathy: a randomized controlled trial. *Am J Sports Med.* 2009;37(3):463–470. <https://doi.org/10.1177/0363546508326983>.
115. Mansur NS, Faloppa F, Bellotti JC, et al. Shock wave therapy associated with eccentric strengthening versus isolated eccentric strengthening for Achilles insertional tendinopathy treatment: a double-blinded randomised clinical trial protocol. *BMJ Open.* 2017;7(1), e013332. <https://doi.org/10.1136/bmjopen-2016-013332>.
116. Wheeler PC, Tattersall C. Novel interventions for recalcitrant Achilles tendinopathy: benefits seen following high-volume image-guided injection or extracorporeal shockwave therapy—a prospective cohort study. *Clin J Sport Med.* 2020;30(1):14–19. <https://doi.org/10.1097/JSM.0000000000000580>.
117. Wheeler PC. Extracorporeal shock wave therapy plus rehabilitation for insertional and noninsertional Achilles tendinopathy shows good results across a range of domains of function. *J Foot Ankle Surg.* 2019;58(4):617–622. <https://doi.org/10.1053/j.jfas.2018.11.005>.
118. Pavone V, Cannavò L, Di Stefano A, Testa G, Costarella L, Sessa G. Low-energy extracorporeal shock-wave therapy in the treatment of chronic insertional Achilles tendinopathy: a case series. *BioMed Res Int.* 2016;2016:7123769. <https://doi.org/10.1155/2016/7123769>.
119. Cutts S, Obi N, Pasapula C, Chan W. Plantar fasciitis. *Ann R Coll Surg Engl.* 2012; 94(8):539–542. <https://doi.org/10.1308/003588412X13171221592456>.
120. Rathleff MS, Thorborg K. 'Load me up, Scotty': mechanotherapy for plantar fasciopathy (formerly known as plantar fasciitis). *Br J Sports Med.* 2015;49(10): 638–639. <https://doi.org/10.1136/bjsports-2014-094562>.
121. Hansen L, Krogh TP, Ellingsen T, Bolvig L, Fredberg U. Long-term prognosis of plantar fasciitis: a 5- to 15-year follow-up study of 174 patients with ultrasound examination. *Orthop J Sports Med.* 2018;6(3): 2325967118757983. <https://doi.org/10.1177/2325967118757983>.
122. Lopes AD, Hespagnol Júnior LC, Yeung SS, Costa LO. What are the main running-related musculoskeletal injuries? A systematic review. *Sports Med.* 2012;42(10): 891–905. <https://doi.org/10.1007/BF03262301>.
123. Monteagudo M, de Albornoz PM, Gutierrez B, Tabuenca J, Álvarez I. Plantar fasciopathy: a current concepts review. *EFORT Open Rev.* 2018;3(8):485–493. <https://doi.org/10.1302/2058-5241.3.170080>.
124. Digiiovanni BF, Nawoczinski DA, Malay DP, et al. Plantar fascia-specific stretching exercise improves outcomes in patients with chronic plantar fasciitis. A prospective clinical trial with two-year follow-up. *J Bone Joint Surg Am.* 2006;88(8):1775–1781. <https://doi.org/10.2106/JBJS.E.01281>.
125. Almubarak A, Foster N. Exercise therapy for plantar heel pain: a systematic review. *Int. J. Exerc. Sci.* 2012;5(3):276–295.
126. Sun J, Gao F, Wang Y, Sun W, Jiang B, Li Z. Extracorporeal shock wave therapy is effective in treating chronic plantar fasciitis: a meta-analysis of RCTs. *Medicine (Baltim).* 2017;96(15), e6621. <https://doi.org/10.1097/MD.0000000000006621>.
127. Li H, Lv H, Lin T. Comparison of efficacy of eight treatments for plantar fasciitis: a network meta-analysis. *J Cell Physiol.* 2018;234(1):860–870. <https://doi.org/10.1002/jcp.26907>.
128. Rompe JD, Furia J, Cacchio A, Schmitz C, Maffulli N. Radial shock wave treatment alone is less efficient than radial shock wave treatment combined with tissue-specific plantar fascia-stretching in patients with chronic plantar heel pain. *Int J Surg.* 2015;24(Pt B):135–142. <https://doi.org/10.1016/j.ijsu.2015.04.082>.
129. Grecco MV, Brech GC, Greve JM. One-year treatment follow-up of plantar fasciitis: radial shockwaves vs. conventional physiotherapy. *Clinics.* 2013;68(8):1089–1095. [https://doi.org/10.6061/clinics/2013\(08\)05](https://doi.org/10.6061/clinics/2013(08)05).
130. Chew KT, Leong D, Lin CY, Lim KK, Tan B. Comparison of autologous conditioned plasma injection, extracorporeal shockwave therapy, and conventional treatment for plantar fasciitis: a randomized trial. *Pharm Manag PM R.* 2013;5(12): 1035–1043. <https://doi.org/10.1016/j.pmjrj.2013.08.590>.
131. Ulusoy A, Cerrahoglu L, Orguc S. Magnetic resonance imaging and clinical outcomes of laser therapy, ultrasound therapy, and extracorporeal shock wave therapy for treatment of plantar fasciitis: a randomized controlled trial. *J Foot Ankle Surg.* 2017;56(4):762–767. <https://doi.org/10.1053/j.jfas.2017.02.013>.
132. Çağlar Okur S, Aydın A. Comparison of extracorporeal shock wave therapy with custom foot orthotics in plantar fasciitis treatment: a prospective randomized one-year follow-up study. *J Musculoskelet Neuronal Interact.* 2019;19(2):178–186.
133. Eslamian F, Shakouri SK, Jahanjoo F, Hajialiloo M, Notghi F. Extra corporeal shock wave therapy versus local corticosteroid injection in the treatment of chronic plantar fasciitis, a single blinded randomized clinical trial. *Pain Med.* 2016;17(9): 1722–1731. <https://doi.org/10.1093/pm/pnw113>.
134. Akınoğlu B, Köse N. A comparison of the acute effects of radial extracorporeal shockwave therapy, ultrasound therapy, and exercise therapy in plantar fasciitis. *J Exerc Rehabil.* 2018;14(2):306–312. <https://doi.org/10.12965/jer.1836048.024>.
135. Vahdatpour B, Mokhtarian A, Raeissadat SA, Dehghan F, Nasr N, Mazaheri M. Enhancement of the effectiveness of extracorporeal shock wave therapy with topical corticosteroid in treatment of chronic plantar fasciitis: a randomized control clinical trial. *Adv Biomed Res.* 2018;7:62. https://doi.org/10.4103/abr.abr_40_17.
136. Takla MKN, Rezk SSR. Clinical effectiveness of multi-wavelength photobiomodulation therapy as an adjunct to extracorporeal shock wave therapy in the management of plantar fasciitis: a randomized controlled trial. *Laser Med Sci.* 2019;34(3):583–593. <https://doi.org/10.1007/s10103-018-2632-4>.
137. Wheeler PC, Tattersall C. Extracorporeal shockwave therapy plus rehabilitation for patients with chronic plantar fasciitis might reduce pain and improve function but still not lead to increased activity: a case-series study with multiple outcome measures. *J Foot Ankle Surg.* 2018;57(2):339–345. <https://doi.org/10.1053/j.jfas.2017.07.001>.
138. Johannsen FE, Herzog RB, Malmgaard-Clausen NM, Hoegberget-Kalish M, Magnusson SP, Kjaer M. Corticosteroid injection is the best treatment in plantar fasciitis if combined with controlled training. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(1):5–12. <https://doi.org/10.1007/s00167-018-5234-6>.
139. Huffer D, Hing W, Newton R, Clair M. Strength training for plantar fasciitis and the intrinsic foot musculature: a systematic review. *Phys Ther Sport.* 2017;24:44–52. <https://doi.org/10.1016/j.pts.2016.08.008>.